

THE MOMENT-SOS HIERARCHY

JEAN B. LASSERRE

Abstract

The Moment-SOS hierarchy initially introduced in optimization in 2000, is based on the theory of the \mathbf{K} -moment problem and its dual counterpart, polynomials that are positive on \mathbf{K} . It turns out that this methodology can be also applied to solve problems with positivity constraints “ $f(\mathbf{x}) \geq 0$ for all $\mathbf{x} \in \mathbf{K}$ ” and/or linear constraints on Borel measures. Such problems can be viewed as specific instances of the “Generalized Problem of Moments” (GPM) whose list of important applications in various domains is endless. We describe this methodology and outline some of its applications in various domains.

1 Introduction

Consider the optimization problem:

$$(1-1) \quad \mathbf{P} : f^* = \inf_{\mathbf{x}} \{ f(\mathbf{x}) : \mathbf{x} \in \Omega \},$$

where f is a polynomial and $\Omega \subset \mathbb{R}^n$ is a basic semi-algebraic set, that is,

$$(1-2) \quad \Omega := \{ \mathbf{x} \in \mathbb{R}^n : g_j(\mathbf{x}) \geq 0, \quad j = 1, \dots, m \},$$

for some polynomials $g_j, j = 1, \dots, m$. Problem \mathbf{P} is a particular case of *Non Linear Programming* (NLP) where the data $(f, g_j, j = 1, \dots, m)$ are *algebraic*, and therefore the whole arsenal of methods of NLP can be used for solving \mathbf{P} . So what is so specific about \mathbf{P} in Equation (1-1)? The answer depends on the meaning of f^* in Equation (1-1).

If one is interested in a *local minimum* only then efficient NLP methods can be used for solving \mathbf{P} . In such methods, the fact that f and g_j 's are polynomials does not help much, that is, this algebraic feature of \mathbf{P} is not really exploited. On the other hand if f^* in Equation (1-1) is understood as the *global minimum* of \mathbf{P} then the picture is totally different. Why? First, to eliminate any ambiguity on the meaning of f^* in Equation (1-1), rewrite Equation (1-1) as:

$$(1-3) \quad \mathbf{P} : f^* = \sup \{ \lambda : f(\mathbf{x}) - \lambda \geq 0, \quad \forall \mathbf{x} \in \Omega \}$$

Research supported by the European Research Council (ERC) through ERC-Advanced Grant # 666981 for the TAMING project.

MSC2010: primary 90C26; secondary 90C22, 90C27, 65K05, 14P10, 44A60.

Keywords: K-Moment problem, positive polynomials, global optimization, semidefinite relaxations.

because then indeed f^* is necessarily the global minimum of \mathbf{P} .

In full generality, most problems [Equation \(1-3\)](#) are very difficult to solve (they are labelled NP-hard in the computational complexity terminology) because:

Given $\lambda \in \mathbb{R}$, checking whether “ $f(\mathbf{x}) - \lambda \geq 0$ for all $\mathbf{x} \in \Omega$ ” is difficult.

Indeed, by nature this positivity constraint is *global* and therefore cannot be handled by standard NLP optimization algorithms which use only local information around a current iterate $\mathbf{x} \in \Omega$. Therefore to compute f^* in [Equation \(1-3\)](#) one needs an efficient tool to handle the positivity constraint “ $f(\mathbf{x}) - \lambda \geq 0$ for all $\mathbf{x} \in \Omega$ ”. Fortunately if the data are algebraic then:

1. Powerful *positivity certificates* from Real Algebraic Geometry (*Positivstellensätze* in german) are available.
2. Some of these positivity certificates have an efficient practical implementation via *Linear Programming* (LP) or *Semidefinite Programming* (SDP). In particular and importantly, testing whether a given polynomial is a sum of squares (SOS) simply reduces to solving a single SDP (which can be done in time polynomial in the input size of the polynomial, up to arbitrary fixed precision).

After the pioneers works of [Shor \[1998\]](#) and [Nesterov \[2000\]](#), [Lasserre \[2000, 2000/01\]](#) and [Parrilo \[2000, 2003\]](#) have been the first to provide a systematic use of these two key ingredients in Control and Optimization, with convergence guarantees. It is also worth mentioning another closely related pioneer work, namely the celebrated SDP-relaxation of [Goemans and Williamson \[1995\]](#) which provides a 0.878 approximation guarantee for MAXCUT, a famous problem in non-convex combinatorial optimization (and probably the simplest one). In fact it is perhaps the first famous example of such a successful application of the powerful SDP convex optimization technique to provide guaranteed good approximations to a notoriously difficult non-convex optimization problem. It turns out that this SDP relaxation is the first relaxation in the Moment-SOS hierarchy (a.k.a. Lasserre hierarchy) when applied to the MAXCUT problem. Since then, this spectacular success story of SDP relaxations has been at the origin of a flourishing research activity in combinatorial optimization and computational complexity. In particular, the study of LP- and SDP-relaxations in hardness of approximation is at the core of a central topic in combinatorial optimization and computational complexity, namely proving/disproving Khot’s famous Unique Games Conjecture¹ (UGC) in Theoretical Computer Science.

Finally, another “definition” of the global optimum f^* of \mathbf{P} reads:

$$(1-4) \quad f^* = \inf_{\mu} \left\{ \int_{\Omega} f d\mu : \mu(\Omega) = 1 \right\}$$

¹For this conjecture and its theoretical and practical implications, S. Khot was awarded the prestigious Nevanlinna prize at the last ICM 2014 in Seoul [Khot \[2014\]](#).

where the ‘inf’ is over all probability measures on Ω . Equivalently, writing f as $\sum_{\alpha} f_{\alpha} \mathbf{x}^{\alpha}$ in the basis of monomials (where $\mathbf{x}^{\alpha} = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$):

$$(1-5) \quad f^* = \inf_{\mathbf{y}} \left\{ \sum_{\alpha} f_{\alpha} y_{\alpha} : \mathbf{y} \in \mathcal{M}(\Omega); \quad y_0 = 1 \right\},$$

where $\mathcal{M}(\Omega) = \{\mathbf{y} = (y_{\alpha})_{\alpha \in \mathbb{N}^n} : \exists \mu \text{ s.t. } y_{\alpha} = \int_{\Omega} \mathbf{x}^{\alpha} d\mu, \forall \alpha \in \mathbb{N}^n\}$, a convex cone. In fact Equation (1-3) is the LP dual of Equation (1-4). In other words standard LP duality between the two formulations Equation (1-4) and Equation (1-3) illustrates the duality between the “ Ω -moment problem” and “polynomials positive on Ω ”.

Problem (1-4) is a very particular instance (and even the simplest instance) of the more general *Generalized Problem of Moments* (GPM):

$$(1-6) \quad \inf_{\mu_1, \dots, \mu_p} \left\{ \sum_{j=1}^p \int_{\Omega_j} f_j d\mu_j : \sum_{j=1}^p \int_{\Omega_j} f_{ij} d\mu_j \geq b_i, i = 1, \dots, s \right\},$$

for some functions $f_{ij} : \mathbb{R}^{n_j} \rightarrow \mathbb{R}, i = 1, \dots, s$, and sets $\Omega_j \subset \mathbb{R}^{n_j}, j = 1, \dots, p$. The GPM is an infinite-dimensional LP with dual:

$$(1-7) \quad \sup_{\lambda_1, \dots, \lambda_s \geq 0} \left\{ \sum_{i=1}^s \lambda_i b_i : f_j - \sum_{i=1}^s \lambda_i f_{ij} \geq 0 \text{ on } \Omega_j, j : 1, \dots, p \right\}.$$

Therefore it should be of no surprise that the Moment-SOS hierarchy, initially developed for global optimization, also applies to solving the GPM. This is particularly interesting as the list of important applications of the GPM is almost endless; see e.g. Landau [1987].

2 The MOMENT-SOS hierarchy in optimization

2.1 Notation, definitions and preliminaries. Let $\mathbb{R}[\mathbf{x}]$ denote the ring of polynomials in the variables $\mathbf{x} = (x_1, \dots, x_n)$ and let $\mathbb{R}[\mathbf{x}]_d$ be the vector space of polynomials of degree at most d (whose dimension is $s(d) := \binom{n+d}{n}$). For every $d \in \mathbb{N}$, let $\mathbb{N}_d^n := \{\alpha \in \mathbb{N}^n : |\alpha| (= \sum_i \alpha_i) \leq d\}$, and let $\mathbf{v}_d(\mathbf{x}) = (\mathbf{x}^{\alpha})_{\alpha \in \mathbb{N}_d^n}$, be the vector of monomials of the canonical basis (\mathbf{x}^{α}) of $\mathbb{R}[\mathbf{x}]_d$. Given a closed set $\mathcal{X} \subseteq \mathbb{R}^n$, let $\mathcal{P}(\mathcal{X}) \subset \mathbb{R}[\mathbf{x}]$ (resp. $\mathcal{P}_d(\mathcal{X}) \subset \mathbb{R}[\mathbf{x}]_d$) be the convex cone of polynomials (resp. polynomials of degree at most $2d$) that are nonnegative on \mathcal{X} . A polynomial $f \in \mathbb{R}[\mathbf{x}]_d$ is written

$$\mathbf{x} \mapsto f(\mathbf{x}) = \sum_{\alpha \in \mathbb{N}^n} f_{\alpha} \mathbf{x}^{\alpha},$$

with vector of coefficients $\mathbf{f} = (f_{\alpha}) \in \mathbb{R}^{s(d)}$ in the canonical basis of monomials $(\mathbf{x}^{\alpha})_{\alpha \in \mathbb{N}^n}$. For real symmetric matrices, let $\langle \mathbf{B}, \mathbf{C} \rangle := \text{trace}(\mathbf{BC})$ while the notation $\mathbf{B} \geq 0$ stands for \mathbf{B} is positive semidefinite (psd) whereas $\mathbf{B} \succ 0$ stands for \mathbf{B} is positive definite (pd).

The Riesz functional. Given a sequence $\mathbf{y} = (y_\alpha)_{\alpha \in \mathbb{N}^n}$, the Riesz functional is the linear mapping $L_{\mathbf{y}} : \mathbb{R}[\mathbf{x}] \rightarrow \mathbb{R}$ defined by:

$$(2-1) \quad f (= \sum_{\alpha} f_{\alpha} \mathbf{x}^{\alpha}) \mapsto L_{\mathbf{y}}(f) = \sum_{\alpha \in \mathbb{N}^n} f_{\alpha} y_{\alpha}.$$

Moment matrix. The *moment* matrix associated with a sequence $\mathbf{y} = (y_{\alpha})_{\alpha \in \mathbb{N}^n}$, is the real symmetric matrix $\mathbf{M}_d(\mathbf{y})$ with rows and columns indexed by \mathbb{N}_d^n , and whose entry (α, β) is just $y_{\alpha+\beta}$, for every $\alpha, \beta \in \mathbb{N}_d^n$. Alternatively, let $\mathbf{v}_d(\mathbf{x}) \in \mathbb{R}^{s(d)}$ be the vector $(\mathbf{x}^{\alpha})_{\alpha \in \mathbb{N}_d^n}$, and define the matrices $(\mathbf{B}_{o,\alpha}) \subset \mathcal{S}^{s(d)}$ by

$$(2-2) \quad \mathbf{v}_d(\mathbf{x}) \mathbf{v}_d(\mathbf{x})^T = \sum_{\alpha \in \mathbb{N}_{2d}^n} \mathbf{B}_{o,\alpha} \mathbf{x}^{\alpha}, \quad \forall \mathbf{x} \in \mathbb{R}^n.$$

Then $\mathbf{M}_d(\mathbf{y}) = \sum_{\alpha \in \mathbb{N}_{2d}^n} y_{\alpha} \mathbf{B}_{o,\alpha}$. If \mathbf{y} has a representing measure μ then $\mathbf{M}_d(\mathbf{y}) \succeq 0$ because $\langle \mathbf{f}, \mathbf{M}_d(\mathbf{y}) \mathbf{f} \rangle = \int f^2 d\mu \geq 0$, for all $f \in \mathbb{R}[\mathbf{x}]_d$.

A measure whose all moments are finite, is *moment determinate* if there is no other measure with same moments. The support of a Borel measure μ on \mathbb{R}^n (denoted $\text{supp}(\mu)$) is the smallest closed set Ω such that $\mu(\mathbb{R}^n \setminus \Omega) = 0$.

Localizing matrix. With \mathbf{y} as above and $g \in \mathbb{R}[\mathbf{x}]$ (with $g(\mathbf{x}) = \sum_{\gamma} g_{\gamma} \mathbf{x}^{\gamma}$), the *localizing* matrix associated with \mathbf{y} and g is the real symmetric matrix $\mathbf{M}_d(g \mathbf{y})$ with rows and columns indexed by \mathbb{N}_d^n , and whose entry (α, β) is just $\sum_{\gamma} g_{\gamma} y_{\alpha+\beta+\gamma}$, for every $\alpha, \beta \in \mathbb{N}_d^n$. Alternatively, let $\mathbf{B}_{g,\alpha} \in \mathcal{S}^{s(d)}$ be defined by:

$$(2-3) \quad g(\mathbf{x}) \mathbf{v}_d(\mathbf{x}) \mathbf{v}_d(\mathbf{x})^T = \sum_{\alpha \in \mathbb{N}_{2d+\text{deg } g}^n} \mathbf{B}_{g,\alpha} \mathbf{x}^{\alpha}, \quad \forall \mathbf{x} \in \mathbb{R}^n.$$

Then $\mathbf{M}_d(g \mathbf{y}) = \sum_{\alpha \in \mathbb{N}_{2d+\text{deg } g}^n} y_{\alpha} \mathbf{B}_{g,\alpha}$. If \mathbf{y} has a representing measure μ whose support is contained in the set $\{\mathbf{x} : g(\mathbf{x}) \geq 0\}$ then $\mathbf{M}_d(g \mathbf{y}) \succeq 0$ for all d because $\langle \mathbf{f}, \mathbf{M}_d(g \mathbf{y}) \mathbf{f} \rangle = \int f^2 g d\mu \geq 0$, for all $f \in \mathbb{R}[\mathbf{x}]_d$.

SOS polynomials and quadratic modules. A polynomial $f \in \mathbb{R}[\mathbf{x}]$ is a Sum-of-Squares (SOS) if there exist $(f_k)_{k=1,\dots,s} \subset \mathbb{R}[\mathbf{x}]$, such that $f(\mathbf{x}) = \sum_{k=1}^s f_k(\mathbf{x})^2$, for all $\mathbf{x} \in \mathbb{R}^n$. Denote by $\Sigma[\mathbf{x}]$ (resp. $\Sigma[\mathbf{x}]_d$) the set of SOS polynomials (resp. SOS polynomials of degree at most $2d$). Of course every SOS polynomial is nonnegative whereas the converse is not true. In addition, checking whether a given polynomial f is nonnegative on \mathbb{R}^n is difficult whereas checking whether f is SOS is much easier and can be done efficiently. Indeed let $f \in \mathbb{R}[\mathbf{x}]_{2d}$ (for f to be SOS its degree must be even), $\mathbf{x} \mapsto f(\mathbf{x}) = \sum_{\alpha \in \mathbb{N}_{2d}^n} f_{\alpha} \mathbf{x}^{\alpha}$. Then f is SOS if and only if there exists a real symmetric matrix $\mathbf{X}^T = \mathbf{X}$ of size $s(d) = \binom{n+d}{n}$, such that:

$$(2-4) \quad \mathbf{X} \succeq 0; \quad f_{\alpha} = \langle \mathbf{X}, \mathbf{B}_{o,\alpha} \rangle, \quad \forall \alpha \in \mathbb{N}_{2d}^n,$$

and this can be checked by solving an SDP.

Next, let $\mathbf{x} \mapsto g_0(\mathbf{x}) := 1$ for all $\mathbf{x} \in \mathbb{R}^n$. With a family $(g_1, \dots, g_m) \subset \mathbb{R}[\mathbf{x}]$ is associated the quadratic module $Q(g) (= Q(g_1, \dots, g_m)) \subset \mathbb{R}[\mathbf{x}]$:

$$(2-5) \quad Q(g) := \left\{ \sum_{j=0}^m \sigma_j g_j : \sigma_j \in \Sigma[\mathbf{x}], j = 0, \dots, m \right\},$$

and its truncated version

$$(2-6) \quad Q_k(g) := \left\{ \sum_{j=0}^m \sigma_j g_j : \sigma_j \in \Sigma[\mathbf{x}]_{k-d_j}, j = 0, \dots, m \right\},$$

where $d_j = \lceil \deg(g_j)/2 \rceil, j = 0, \dots, m$.

Definition 1. The quadratic module $Q(g)$ associated with Ω in Equation (1-2) is said to be Archimedean if there exists $M > 0$ such that the quadratic polynomial $\mathbf{x} \mapsto M - \|\mathbf{x}\|^2$ belongs to $Q(g)$ (i.e., belongs to $Q_k(g)$ for some k).

If $Q(g)$ is Archimedean then necessarily Ω is compact but the reverse is not true. The Archimedean condition (which depends on the representation of Ω) can be seen as an algebraic certificate that Ω is compact. For more details on the above notions of moment and localizing matrix, quadratic module, as well as their use in potential applications, the interested reader is referred to Lasserre [2010], Laurent [2009], Schmüdgen [2017].

2.2 Two certificates of positivity (Positivstellensätze). Below we describe two particular certificates of positivity which are important because they provide the theoretical justification behind the so-called SDP- and LP-relaxations for global optimization.

Theorem 2.1 (Putinar [1993]). Let $\Omega \subset \mathbb{R}^n$ be as in Equation (1-2) and assume that $Q(g)$ is Archimedean.

- (a) If a polynomial $f \in \mathbb{R}[\mathbf{x}]$ is (strictly) positive on Ω then $f \in Q(g)$.
- (b) A sequence $\mathbf{y} = (y_\alpha)_{\alpha \in \mathbb{N}^n} \subset \mathbb{R}$ has a representing Borel measure on Ω if and only if $L_{\mathbf{y}}(f^2 g_j) \geq 0$ for all $f \in \mathbb{R}[\mathbf{x}]$, and all $j = 0, \dots, m$. Equivalently, if and only if $\mathbf{M}_d(\mathbf{y} g_j) \geq 0$ for all $j = 0, \dots, m, d \in \mathbb{N}$.

There exists another certificate of positivity which does not use SOS.

Theorem 2.2 (Krivine [1964a], Krivine [1964b], and Vasilescu [2003]). Let $\Omega \subset \mathbb{R}^n$ as in Equation (1-2) be compact and such that (possibly after scaling) $0 \leq g_j(\mathbf{x}) \leq 1$ for all $\mathbf{x} \in \Omega, j = 1, \dots, m$. Assume also that $[1, g_1, \dots, g_m]$ generates $\mathbb{R}[\mathbf{x}]$.

- (a) If a polynomial $f \in \mathbb{R}[\mathbf{x}]$ is (strictly) positive on Ω then

$$(2-7) \quad f(\mathbf{x}) = \sum_{\alpha, \beta \in \mathbb{N}^n} c_{\alpha, \beta} \prod_{j=1}^m g_j(\mathbf{x})^{\alpha_j} (1 - g_j(\mathbf{x}))^{\beta_j},$$

for finitely many positive coefficients $(c_{\alpha,\beta})_{\alpha,\beta \in \mathbb{N}^m}$.

(b) A sequence $\mathbf{y} = (y_\alpha)_{\alpha \in \mathbb{N}^n} \subset \mathbb{R}$ has a representing Borel measure on Ω if and only if $L_{\mathbf{y}} \left(\prod_{j=1}^m g_j(\mathbf{x})^{\alpha_j} (1 - g_j(\mathbf{x}))^{\beta_j} \right) \geq 0$ for all $\alpha, \beta \in \mathbb{N}^m$.

The two facets (a) and (b) of Theorem 2.1 and Theorem 2.2 illustrate the duality between polynomials positive on Ω (in (a)) and the Ω -moment problem (in (b)). In addition to their mathematical interest, both Theorem 2.1(a) and Theorem 2.2(a) have another distinguishing feature. They both have a practical implementation. Testing whether $f \in \mathbb{R}[\mathbf{x}]_d$ is in $Q(g)_k$ is just solving a single SDP, whereas testing whether f can be written as in Equation (2-7) with $\sum_{i=1}^m \alpha_i + \beta_i \leq k$, is just solving a single Linear Program (LP).

2.3 The Moment-SOS hierarchy. The Moment-SOS hierarchy is a numerical scheme based on Putinar’s theorem. In a nutshell it consists of replacing the intractable positivity constraint “ $f(\mathbf{x}) \geq 0$ for all $\mathbf{x} \in \Omega$ ” with Putinar’s positivity certificate $f \in Q_d(g)$ of Theorem 2.1(a), i.e., with a fixed degree bound on the SOS weights (σ_j) in Equation (2-6). By duality, it consists of replacing the intractable constraint $\mathbf{y} \in \mathcal{M}(\Omega)$ with the necessary conditions $\mathbf{M}_d(g_j \mathbf{y}) \geq 0, j = 0, \dots, m$, of Theorem 2.1(b) for a fixed d . This results in solving an SDP which provides a lower bound on the global minimum. By allowing the degree bound d to increase, one obtains a hierarchy of SDPs (of increasing size) which provides a monotone non-decreasing sequence of lower bounds. A similar strategy based on Krivine-Stengle-Vasilescu positivity certificate (Equation (2-7)) is also possible and yields a hierarchy of LP (instead of SDPs). However even though one would prefer to solve LPs rather than SDPs, the latter Moment-LP hierarchy has several serious drawbacks (some explained in e.g. Lasserre [2015a, 2002b]), and therefore we only describe the Moment-SOS hierarchy.

Recall problem **P** in Equation (1-1) or equivalently in Equation (1-3) and Equation (1-4), where $\Omega \subset \mathbb{R}^n$ is the basic semi-algebraic set defined in Equation (1-2).

The Moment-SOS hierarchy. Consider the sequence of semidefinite programs $(\mathbf{Q}_d)_{d \in \mathbb{N}}$ with $d \geq \hat{d} := \max[\deg(f), \max_j \deg(g_j)]$:

$$(2-8) \quad \mathbf{Q}_d : \rho_d = \inf_{\mathbf{y}} \{ L_{\mathbf{y}}(f) : y_0 = 1; \mathbf{M}_d(g_j \mathbf{y}) \geq 0, \quad 0 \leq j \leq m \}$$

(where $\mathbf{y} = (y_\alpha)_{\alpha \in \mathbb{N}_{2d}^n}$)², with associated sequence of their SDP duals:

$$(2-9) \quad \mathbf{Q}_d^* : \rho_d^* = \sup_{\lambda, \sigma_j} \{ \lambda : f - \lambda = \sum_{j=0}^m \sigma_j g_j; \sigma_j \in \Sigma[\mathbf{x}]_{d-d_j}, \quad 0 \leq j \leq m \}$$

(where $d_j = \lceil (\deg g_j)/2 \rceil$). By standard weak duality in optimization $\rho_d^* \leq \rho_d$ for every $d \geq \hat{d}$. The sequence $(\mathbf{Q}_d)_{d \in \mathbb{N}}$ forms a hierarchy of SDP-relaxations of **P**

²In Theoretical Computer Science, \mathbf{y} is called a sequence of “pseudo-moments”.

because $\rho_d \leq f^*$ and $\rho_d \leq \rho_{d+1}$ for all $d \geq \hat{d}$. Indeed for each $d \geq \hat{d}$, the constraints of \mathbf{Q}_d consider only necessary conditions for \mathbf{y} to be the moment sequence (up to order $2d$) of a probability measure on Ω (cf. [Theorem 2.1\(b\)](#)) and therefore \mathbf{Q}_d is a relaxation of [Equation \(1-5\)](#).

By duality, the sequence $(\mathbf{Q}_d^*)_{d \in \mathbb{N}}$ forms a hierarchy of SDP-strengthenings of [Equation \(1-3\)](#). Indeed in [Equation \(2-9\)](#) one has replaced the intractable positivity constraint of [Equation \(1-3\)](#) by the (stronger) Putinar’s positivity certificate with degree bound $2d - 2d_j$ on the SOS weights σ_j ’s.

Theorem 2.3 ([Lasserre \[2000, 2000/01\]](#)). *Let Ω in [Equation \(1-2\)](#) be compact and assume that its associated quadratic module $Q(g)$ is Archimedean. Then:*

(i) *As $d \rightarrow \infty$, the monotone non-decreasing sequence $(\rho_d)_{d \in \mathbb{N}}$ (resp. $(\rho_d^*)_{d \in \mathbb{N}}$) of optimal values of the hierarchy ([Equation \(2-8\)](#)) (resp. [Equation \(2-9\)](#)) converges to the global optimum f^* of \mathbf{P} .*

(ii) *Moreover, let $\mathbf{y}^d = (y_\alpha^d)_{\alpha \in \mathbb{N}_{2d}^n}$ be an optimal solution of \mathbf{Q}_d in [Equation \(2-8\)](#), and let $s = \max_j d_j$ (recall that $d_j = \lceil (\deg g_j)/2 \rceil$). If*

$$(2-10) \quad \text{rank } \mathbf{M}_d(\mathbf{y}^d) = \text{rank } \mathbf{M}_{d-s}(\mathbf{y}^d) (=: t)$$

then $\rho_d = f^$ and there are t global minimizers $\mathbf{x}_j^* \in \Omega$, $j = 1, \dots, t$, that can be “extracted” from \mathbf{y}^d by a linear algebra routine.*

The sequence of SDP-relaxations (\mathbf{Q}_d) , $d \geq \hat{d}$, and the rank test ([Equation \(2-10\)](#)) to extract global minimizers, are implemented in the GloptiPoly software [Henrion, Lasserre, and Löfberg \[2009\]](#).

Finite convergence and a global optimality certificate. After being introduced in [Lasserre \[2000\]](#), in many numerical experiments it was observed that typically, finite convergence takes place, that is, $f^* = \rho_d$ for some (usually small) d . In fact there is a rationale behind this empirical observation.

Theorem 2.4 ([Nie \[2014a\]](#)). *Let \mathbf{P} be as in [Equation \(1-3\)](#) where Ω in [Equation \(1-2\)](#) is compact and its associated quadratic module is Archimedean. Suppose that at each global minimizer $\mathbf{x}^* \in \Omega$:*

- *The gradients $(\nabla g_j(\mathbf{x}^*))_{j=1, \dots, m}$ are linearly independent. (This implies existence of nonnegative Lagrange-KKT multipliers λ_j^* , $j \leq m$, such that $\nabla f(\mathbf{x}^*) - \sum_{j=1}^m \lambda_j^* \nabla g_j(\mathbf{x}^*) = 0$ and $\lambda_j^* g_j(\mathbf{x}^*) = 0$ for all $j \leq m$.)*
- *Strict complementarity holds, that is, $g_j(\mathbf{x}^*) = 0 \Rightarrow \lambda_j^* > 0$.*
- *Second-order sufficiency condition holds, i.e.,*

$$\langle \mathbf{u}, \nabla_{\mathbf{x}}^2 (f(\mathbf{x}^*) - \sum_{j=1}^m \lambda_j^* g_j(\mathbf{x}^*)) \mathbf{u} \rangle > 0,$$

for all $0 \neq \mathbf{u} \in \nabla(f(\mathbf{x}^) - \sum_{j=1}^m \lambda_j^* g_j(\mathbf{x}^*))^\perp$.*

Then $f - f^* \in Q(g)$, i.e., there exists d^* and SOS multipliers $\sigma_j^* \in \Sigma[\mathbf{x}]_{d^*-d_j}$, $j = 0, \dots, m$, such that:

$$(2-11) \quad f(\mathbf{x}) - f^* = \sigma_0^*(\mathbf{x}) + \sum_{j=1}^m \sigma_j^*(\mathbf{x}) g_j(\mathbf{x}).$$

With Equation (2-11), Theorem 2.4 provides a *certificate of global optimality* in polynomial optimization, and to the best of our knowledge, the first at this level of generality. Next, observe that $\mathbf{x}^* \in \Omega$ is a global unconstrained minimizer of the *extended Lagrangian polynomial* $f - f^* - \sum_{j=1}^m \sigma_j^* g_j$, and therefore Theorem 2.4 is the analogue for *non-convex* polynomial optimization of the Karush-Kuhn-Tucker (KKT) optimality conditions *in the convex case*. Indeed in the convex case, any local minimizer is global and is also a global unconstrained minimizer of the Lagrangian $f - f^* - \sum_{j=1}^m \lambda_j^* g_j$.

Also interestingly, whenever the SOS weight σ_j^* in Equation (2-11) is non trivial, it testifies that the constraint $g_j(\mathbf{x}) \geq 0$ is important for \mathbf{P} even if it is not active at \mathbf{x}^* (meaning that if $g_j \geq 0$ is deleted from \mathbf{P} then the new global optimum decreases strictly). The multiplier λ_j^* plays the same role in the KKT-optimality conditions *only in the convex case*. See Lasserre [2015a] for a detailed discussion.

Finite convergence of the Moment-SOS-hierarchies (Equations (2-8) and (2-9)) is an immediate consequence of Theorem 2.4. Indeed by Equation (2-11) $(f^*, \sigma_0^*, \dots, \sigma_m^*)$ is a feasible solution of $\mathbf{Q}_{d^*}^*$ with value $f^* \leq \rho_d^* \leq f^*$ (hence $\rho_d^* = \rho_d = f^*$).

Genericity: Importantly, as proved in Nie [2014a], the conditions in Theorem 2.4 are *generic*. By this we mean the following: Consider the class $\mathcal{P}(t, m)$ of optimization problems \mathbf{P} with data (f, g_1, \dots, g_m) of degree bounded by t , and with nonempty compact feasible set Ω . Such a problem \mathbf{P} is a “point” in the space $\mathbb{R}^{(m+1)s(t)}$ of coordinates of (f, g_1, \dots, g_m) . Then the “good” problems \mathbf{P} are points in a Zariski open set. Moreover, generically the rank test (Equation (2-10)) is also satisfied at an optimal solution of Equation (2-8) (for some d); for more details see Nie [2013].

Computational complexity: Each relaxation \mathbf{Q}_d in Equation (2-8) is a semidefinite program with $s(2d) = \binom{n+2d}{n}$ variables (y_α) , and a psd constraint $\mathbf{M}_d(\mathbf{y}) \succeq 0$ of size $s(d)$. Therefore solving \mathbf{Q}_d in its canonical form Equation (2-8) is quite expensive in terms of computational burden, especially when using interior-point methods. Therefore its brute force application is limited to small to medium size problems.

Exploiting sparsity: Fortunately many large scale problems exhibit a structured sparsity pattern (e.g., each polynomial g_j is concerned with a few variables only, and the objective function f is a sum $\sum_i f_i$ where each f_i is also concerned with a few variables only). Then Waki, Kim, Kojima, and Muramatsu [2006] have proposed a sparsity-adapted hierarchy of SDP-relaxations which can handle problems \mathbf{P} with thousands variables. In addition, if the sparsity pattern satisfies a certain condition then convergence of this sparsity-adapted hierarchy is also guaranteed like in the dense case Lasserre [2006]. Successful applications of this strategy can be found in e.g. Laumond, Mansard, and Lasserre [2017a] in Control (systems identification) and in Molzahn and Hiskens [2015]

for solving (large scale) Optimum Power Flow problems (OPF is an important problem encountered in the management of energy networks).

2.4 Discussion. We claim that the Moment-SOS hierarchy and its rationale [Theorem 2.4](#), unify convex, non-convex (continuous), and discrete (polynomial) Optimization. Indeed in the description of \mathbf{P} we do not pay attention to what particular class of problems \mathbf{P} belongs to. This is in sharp contrast to the usual common practice in (local) optimization where several classes of problems have their own tailored favorite class of algorithms. For instance, problems are not treated the same if equality constraints appear, and/or if boolean (or discrete variables) are present, etc. Here a boolean variable x_i is modeled by the quadratic equality constraint $x_i^2 = x_i$. So it is reasonable to speculate that this lack of specialization could be a handicap for the moment-SOS hierarchy.

But this is not so. For instance for the sub-class of convex³ problems \mathbf{P} where f and $(-g_j)_{j=1,\dots,m}$ are SOS-convex⁴ polynomials, finite convergence takes place at the first step of the hierarchy. In other words, the SOS hierarchy somehow “recognizes” this class of easy problems [Lasserre \[2015a\]](#). In the same time, for a large class of 0/1 combinatorial optimization problems on graphs, the Moment-SOS hierarchy has been shown to provide the tightest upper bounds when compared to the class of *lift-and-project* methods, and has now become a central tool to analyze hardness of approximations in combinatorial optimization. For more details the interested reader is referred to e.g. [Lasserre \[2002b\]](#), [Laurent \[2003\]](#), [Barak and Steurer \[2014\]](#), [Khot \[2010, 2014\]](#) and the many references therein.

3 The Moment-SOS hierarchy outside optimization

3.1 A general framework for the Moment-SOS hierarchy. Let $\Omega_i \subset \mathbb{R}^{n_i}$ be a finite family of compact sets, $\mathcal{M}(\Omega_i)$ (resp. $\mathcal{C}(\Omega_i)$) be the space of finite Borel signed measures (resp. continuous functions) on Ω_i , $i = 0, 1, \dots, s$, and let \mathbf{T} be a continuous linear mapping with adjoint \mathbf{T}^* :

$$\begin{aligned} \mathbf{T} : \mathcal{M}(\Omega_1) \times \dots \times \mathcal{M}(\Omega_s) &\rightarrow \mathcal{M}(\Omega_0) \\ \mathcal{C}(\Omega_1) \times \dots \times \mathcal{C}(\Omega_s) &\leftarrow \mathcal{C}(\Omega_0) : \mathbf{T}^* \end{aligned}$$

Let $\phi := (\phi_1, \dots, \phi_s)$ and let $\phi_i \geq 0$ stand for ϕ_i is a positive measure. Then consider the general framework:

$$(3-1) \quad \rho = \inf_{\phi \geq 0} \left\{ \sum_{i=1}^s \langle f_i, \phi_i \rangle : \mathbf{T}(\phi) = \lambda; \sum_{i=1}^s \langle f_{ij}, \phi_i \rangle \geq b_j, j \in J \right\},$$

³Convex problems \mathbf{P} where f and $(-g_j)_{j=1,\dots,m}$ are convex, are considered “easy” and can be solved efficiently.

⁴A polynomial $f \in \mathbb{R}[\mathbf{x}]$ is SOS-convex if its Hessian $\nabla^2 f$ is a SOS matrix-polynomial, i.e., $\nabla f^2(\mathbf{x}) = \mathbf{L}(\mathbf{x})\mathbf{L}(\mathbf{x})^T$ for some matrix-polynomial $\mathbf{L} \in \mathbb{R}[\mathbf{x}]^{n \times p}$.

where J is a finite or countable set, $\mathbf{b} = (b_j)$ is given, $\lambda \in \mathcal{M}(\Omega_0)$ is a given measure, $(f_{ij})_{j \in J}, i = 1, \dots, s$, are given polynomials, and $\langle \cdot, \cdot \rangle$ is the duality bracket between $\mathcal{C}(\Omega_i)$ and $\mathcal{M}(\Omega_i)$ ($\langle h, \phi_i \rangle = \int_{\Omega_i} h d\phi_i, i = 1, \dots, s$).

As we will see, this general framework is quite rich as it encompasses a lot of important applications in many different fields. In fact Problem (3-1) is equivalent to the Generalized Problem of Moments (GPM):

$$(3-2) \quad \rho = \inf_{\phi \geq 0} \left\{ \sum_{i=1}^s \langle f_i, \phi_i \rangle : \langle \mathbf{T}^* p_k, \phi \rangle = \langle p_k, \lambda \rangle, \quad k = 0, 1, \dots, \right. \\ \left. \sum_{i=1}^s \langle f_{ij}, \phi_i \rangle \geq b_j, \quad j \in J \right\},$$

where the family $(p_k)_{k=0, \dots}$ is dense in $\mathcal{C}(\Omega_0)$ (e.g. a basis of $\mathbb{R}[x_1, \dots, x_{n_0}]$).

The Moment-SOS hierarchy can also be applied to help solve the Generalized Problem of Moments (GPM) (Equation (3-2)) or its dual :

$$(3-3) \quad \rho^* = \sup_{(\theta_j \geq 0, \square)} \left\{ \sum_k \gamma_k \langle p_k, \lambda \rangle + \langle \theta, \mathbf{b} \rangle : \right. \\ \left. \text{s.t. } f_i - \sum_k \gamma_k (\mathbf{T}^* p_k)_i - \sum_{j \in J} \theta_j f_{ij} \geq 0 \text{ on } \Omega_i \text{ for all } i \right\},$$

where the unknown $\gamma = (\gamma_k)_{k \in \mathbb{N}}$ is a finite sequence.

3.2 A hierarchy of SDP-relaxations. Let

$$(3-4) \quad \Omega_i := \{ \mathbf{x} \in \mathbb{R}^{n_i} : g_{i,\ell}(\mathbf{x}) \geq 0, i = 1, \dots, m_i \}, \quad i = 1, \dots, s,$$

for some polynomials $(g_{i,\ell}) \subset \mathbb{R}[x_1, \dots, x_{n_i}], \ell = 1, \dots, m_i$. Let $d_{i,\ell} = \lceil \deg(g_{i,\ell})/2 \rceil$ and $\hat{d} := \max_{i,j,\ell} [\deg(f_i), \deg(f_{ij}), \deg(g_{i,\ell})]$. To solve Equation (3-2), define the ‘‘moment’’ sequences $\mathbf{y}_i = (y_{i,\alpha}), \alpha \in \mathbb{N}^{n_i}, i = 1, \dots, s$, and with $d \in \mathbb{N}$, define $\Gamma_d := \{ p_k : \deg(\mathbf{T}^* p_k)_i \leq 2d, i = 1, \dots, s \}$. Consider the hierarchy of semidefinite programs indexed by $\hat{d} \leq d \in \mathbb{N}$:

$$(3-5) \quad \rho_d = \inf_{(\mathbf{y}_i)} \left\{ \sum_{i=1}^s L_{\mathbf{y}_i}(f_i) : \sum_{i=1}^s L_{\mathbf{y}_i}((\mathbf{T}^* p_k)_i) = \langle p_k, \lambda \rangle, \quad p_k \in \Gamma_d \right. \\ \left. \sum_{i=1}^s L_{\mathbf{y}_i}(f_{ij}) \geq b_j, \quad j \in J_d \right. \\ \left. \mathbf{M}_d(\mathbf{y}_i), \mathbf{M}_{d-d_\ell}(g_{i\ell} \mathbf{y}_i) \geq 0, \quad \ell \leq m_i; i \leq s \right\},$$

where $J_d \subset J$ is finite $\bigcup_{d \in \mathbb{N}} J_d = J$. Its dual SDP-hierarchy reads:

$$(3-6) \quad \rho_d^* = \sup_{(\theta_j \geq 0, \gamma_k)} \left\{ \sum_{p_k \in \Gamma_d} \gamma_k \langle p_k, \lambda \rangle + \langle \theta, \mathbf{b} \rangle : \right. \\ \left. \text{s.t. } f_i - \sum_{p_k \in \Gamma_d} \gamma_k (\mathbf{T}^* p_k)_i - \sum_{j \in J} \theta_j f_{ij} = \sum_{\ell=0}^{m_i} \sigma_{i,\ell} g_{i,\ell} \right. \\ \left. \sigma_{i,\ell} \in \Sigma[x_1, \dots, x_{n_i}]_{d-d_{i,\ell}}; i = 1, \dots, s \right\},$$

As each Ω_i is compact, for technical reasons and with no loss of generality, in the sequel we may and will assume that for every $i = 1, \dots, s$, $g_{i,0}(\mathbf{x}) = M_i - \|\mathbf{x}\|^2$, where $M_i > 0$ is sufficiently large.

Theorem 3.1. *Assume that $\rho > -\infty$ and that for every $i = 1, \dots, s$, $f_{i0} = 1$. Then for every $d \geq \hat{d}$, Equation (3-5) has an optimal solution, and $\lim_{d \rightarrow \infty} \rho_d = \rho$.*

3.3 Example In Probability and Computational Geometry.

Bounds on measures with moment conditions. Let Z be a random vector with values in a compact semi-algebraic set $\Omega_1 \subset \mathbb{R}^n$. Its distribution λ on Ω_1 is unknown but some of its moments $\int \mathbf{x}^\alpha d\lambda = b_\alpha$, $\alpha \in \Gamma \subset \mathbb{N}^n$, are known ($b_0 = 1$). Given a basic semi-algebraic set $\Omega_2 \subset \Omega_1$ we want to compute (or approximate as closely as desired) the best upper bound on $\text{Prob}(Z \in \Omega_2)$. This problem reduces to solving the GPM:

$$(3-7) \quad \rho = \sup_{\phi_1, \phi_2 \geq 0} \{ \langle 1, \phi_2 \rangle : \langle \mathbf{x}^\alpha, \phi_1 \rangle + \langle \mathbf{x}^\alpha, \phi_2 \rangle = b_\alpha, \alpha \in \Gamma; \phi_i \in \mathcal{M}(\Omega_i), i = 1, 2 \},$$

With Ω_1 and Ω_2 as in Equation (3-4) one may compute upper bounds on ρ by solving the Moment-SOS hierarchy (Equation (3-5)) adapted to problem (Equation (3-7)). Under the assumptions of Theorem 3.1, the resulting sequence $(\rho_d)_{d \in \mathbb{N}}$ converges to ρ as $d \rightarrow \infty$; for more details the interested reader is referred to Lasserre [2002a].

Lebesgue & Gaussian measures of semi-algebraic sets. Let $\Omega_2 \subset \mathbb{R}^n$ be compact. The goal is to compute (or approximate as closely as desired) the Lebesgue measure $\lambda(\Omega_2)$ of Ω_2 . Then take $\Omega_1 \supset \Omega_2$ be a simple set, e.g. an ellipsoid or a box (in fact any set such that one knows all moments $(b_\alpha)_{\alpha \in \mathbb{N}^n}$ of the Lebesgue measure on Ω_1). Then:

$$(3-8) \quad \lambda(\Omega_2) = \sup_{\phi_1, \phi_2 \geq 0} \{ \langle 1, \phi_2 \rangle : \langle \mathbf{x}^\alpha, \phi_1 \rangle + \langle \mathbf{x}^\alpha, \phi_2 \rangle = b_\alpha, \alpha \in \mathbb{N}^n; \phi_i \in \mathcal{M}(\Omega_i), i = 1, 2 \}.$$

Problem (3-8) is very similar to (3-7) except that we now have countably many moment constraints ($\Gamma = \mathbb{N}^n$). Again, with Ω_2 and Ω_1 as in Equation (3-4) one may compute upper bounds on $\lambda(\Omega_2)$ by solving the Moment-SOS hierarchy (Equation (3-5)) adapted to problem (3-8). Under the assumptions of Theorem 3.1, the resulting monotone non-increasing sequence $(\rho_d)_{d \in \mathbb{N}}$ converges to $\lambda(\Omega_2)$ from above as $d \rightarrow \infty$. The convergence $\rho_d \rightarrow \lambda(\Omega_2)$ is slow because of a Gibb's phenomenon⁵. Indeed the semidefinite program (Equation (3-6)) reads:

$$\rho_d^* = \inf_{p \in \mathbb{R}[\mathbf{x}]_{2d}} \left\{ \int_{\Omega_1} p d\lambda : p \geq 1 \text{ on } \Omega_2; \quad p \geq 0 \text{ on } \Omega_1 \right\},$$

⁵The Gibbs' phenomenon appears at a jump discontinuity when one approximates a piecewise C^1 function with a continuous function, e.g., by its Fourier series.

i.e., as $\rightarrow \infty$ one tries to approximate the discontinuous function $\mathbf{x} \mapsto 1_{\Omega_2}(\mathbf{x})$ by polynomials of increasing degrees. Fortunately there are several ways to accelerate the convergence, e.g. as in [Henrion, Lasserre, and Savorgnan \[2009\]](#) (but loosing the monotonicity) or in [Lasserre \[2017\]](#) (preserving monotonicity) by including in [Equation \(3-5\)](#) additional constraints on \mathbf{y}_2 coming from an application of Stokes' theorem.

For the **Gaussian measure** λ we need and may take $\Omega_1 = \mathbb{R}^n$ and Ω_2 is not necessarily compact. Although both Ω_1 and Ω_2 are allowed to be non-compact, the Moment-SOS hierarchy ([Equation \(3-5\)](#)) still converges, i.e., $\rho_d \rightarrow \lambda(\Omega_2)$ as $d \rightarrow \infty$. This is because the moments of λ satisfy the generalized Carleman's condition

$$(3-9) \quad \sum_{k=1}^{\infty} \left(\int_{\mathbb{R}^n} x_i^{2k} d\lambda \right)^{-1/2k} = +\infty, \quad i = 1, \dots, n,$$

which imposes implicit constraints on \mathbf{y}_1 and \mathbf{y}_2 in [Equation \(3-5\)](#), strong enough to guarantee $\rho_d \rightarrow \lambda(\Omega_2)$ as $d \rightarrow \infty$. For more details see [Lasserre \[ibid.\]](#). This deterministic approach is computationally demanding and should be seen as complementary to brute force Monte-Carlo methods that provide only an estimate (but can handle larger size problems).

3.4 In signal processing and interpolation. In this application, a signal is identified with an atomic signed measure ϕ supported on few atoms $(\mathbf{x}_k)_{k=1,\dots,s} \subset \Omega$, i.e., $\phi = \sum_{k=1}^s \theta_k \delta_{\mathbf{x}_k}$, for some weights $(\theta_k)_{k=1,\dots,s}$.

Super-Resolution. The goal of Super-Resolution is to reconstruct the unknown measure ϕ (the signal) from a few measurements only, when those measurements are the moments $(b_\alpha)_{\alpha \in \mathbb{N}_t^n}$ of ϕ , up to order t (fixed). One way to proceed is to solve the infinite-dimensional program:

$$(3-10) \quad \rho = \inf_{\phi} \{ \|\phi\|_{TV} : \int \mathbf{x}^\alpha d\phi = b_\alpha, \quad \alpha \in \mathbb{N}_t^n \},$$

where the inf is over the finite signed Borel measures on Ω , and $\|\phi\|_{TV} = |\phi|(\Omega)$ (with $|\phi|$ being the total variation of ϕ). Equivalently:

$$(3-11) \quad \rho = \inf_{\phi^+, \phi^- \geq 0} \{ \langle 1, \phi^+ + \phi^- \rangle : \langle \mathbf{x}^\alpha, \phi^+ - \phi^- \rangle = b_\alpha, \quad \alpha \in \mathbb{N}_t^n \},$$

which is an instance of the GPM with dual:

$$(3-12) \quad \rho^* = \sup_{p \in \mathbb{R}[\mathbf{x}]_t} \left\{ \sum_{\alpha \in \mathbb{N}_t^n} p_\alpha b_\alpha : \|p\|_\infty \leq 1 \right\},$$

where $\|p\|_\infty = \sup\{|p(\mathbf{x})| : \mathbf{x} \in \Omega\}$. In this case, the Moment-SOS hierarchy ([Equation \(3-5\)](#)) with $d \geq \hat{d} := \lceil t/2 \rceil$, reads:

$$(3-13) \quad \begin{aligned} \rho_d = \inf_{\mathbf{y}^+, \mathbf{y}^-} \{ & y_0^+ + y_0^- : y_\alpha^+ - y_\alpha^- = b_\alpha, \quad \alpha \in \mathbb{N}_t^n \\ & \mathbf{M}_d(\mathbf{y}^\pm) \geq 0; \mathbf{M}_d(g_\ell \mathbf{y}^\pm) \geq 0, \quad \ell = 1, \dots, m \}, \end{aligned}$$

where $\Omega = \{\mathbf{x} : g_\ell(\mathbf{x}) \geq 0, \ell = 1, \dots, m\}$.

In the case where Ω is the torus $\mathbb{T} \subset \mathbb{C}$, Candès and Fernandez-Granda [2014] showed that if $\delta > 2/f_c$ (where δ is the minimal distance between the atoms of ϕ , and f_c is the number of measurements) then Equation (3-10) has a unique solution and one may recover ϕ exactly by solving the single semidefinite program (Equation (3-10)) with $d = \lceil t/2 \rceil$. The dual (Equation (3-12)) has an optimal solution p^* (a trigonometric polynomial) and the support of ϕ^+ (resp. ϕ^-) consists of the atoms $\mathbf{z} \in \mathbb{T}$ of ϕ such that $p^*(\mathbf{z}) = 1$ (resp. $p^*(\mathbf{z}) = -1$). In addition, this procedure is more robust to noise in the measurements than Prony’s method; on the other hand, the latter requires less measurements and no separation condition on the atoms.

In the general multivariate case treated in De Castro, Gamboa, Henrion, and Lasserre [2017] one now needs to solve the Moment-SOS hierarchy (Equation (3-11)) for $d = \hat{d}, \dots$ (instead of a single SDP in the univariate case). However since the moment constraints of Equation (3-11) are finitely many, exact recovery (i.e. finite convergence of the Moment-SOS hierarchy (Equation (3-13))) is possible (usually with a few measurements only). This is indeed what has been observed in all numerical experiments of De Castro, Gamboa, Henrion, and Lasserre [ibid.], and in all cases with significantly less measurements than the theoretical bound (of a tensorized version of the univariate case).

In fact, the rank condition (Equation (2-10)) is always satisfied at an optimal solution $(\mathbf{y}^+, \mathbf{y}^-)$ at some step d of the hierarchy (Equation (3-13)), and so the atoms of ϕ^+ and ϕ^- are extracted via a simple linear algebra routine (as for global optimization). Nie’s genericity result Nie [2013] should provide a rationale which explains why the rank condition (Equation (2-10)) is satisfied in all examples.

Sparse interpolation. Here the goal is to recover an unknown (black-box) polynomial $p \in \mathbb{R}[\mathbf{x}]_t$ through a few evaluations of p only. In Josz, Lasserre, and Mourrain [2017] we have shown that this problem is in fact a particular case of Super-Resolution (and even discrete Super-Resolution) on the torus $\mathbb{T}^n \subset \mathbb{C}^n$. Indeed let $\mathbf{z}_0 \in \mathbb{T}^n$ be fixed, arbitrary. Then with $\beta \in \mathbb{N}^n$, notice that

$$\begin{aligned} p(\mathbf{z}_0^\beta) &= \sum_{\alpha \in \mathbb{N}_d^n} p_\alpha (z_{01}^{\beta_1} \cdots z_{0n}^{\beta_n})^\alpha = \sum_{\alpha \in \mathbb{N}_d^n} p_\alpha (z_{01}^{\alpha_1} \cdots z_{0n}^{\alpha_n})^\beta \\ &= \int_{\mathbb{T}^n} \mathbf{z}^\beta d \left(\sum_{\alpha \in \mathbb{N}_d^n} p_\alpha \delta_{\mathbf{z}_0^\alpha} \right) = \int_{\mathbb{T}^n} \mathbf{z}^\beta d\phi. \end{aligned}$$

In other words, one may identify the polynomial p with an atomic signed Borel measure ϕ on \mathbb{T}^n supported on finitely many atoms $(\mathbf{z}_0^\alpha)_{\alpha \in \mathbb{N}_d^n}$ with associated weights $(p_\alpha)_{\alpha \in \mathbb{N}_d^n}$.

Therefore, if the evaluations of the black-box polynomial p are done at a few “powers” $(\mathbf{z}_0^\beta), \beta \in \mathbb{N}^n$, of an arbitrary point $\mathbf{z}_0 \in \mathbb{T}^n$, then the sparse interpolation problem is equivalent to recovering an unknown atomic signed Borel measure ϕ on \mathbb{T}^n from knowledge of a few moments, that is, the Super-Resolution problem that we have just

described above. Hence one may recover p by solving the Moment-SOS hierarchy (Equation (3-13)) for which finite convergence usually occurs fast. For more details see Jozs, Lasserre, and Mourrain [2017].

3.5 In Control & Optimal Control. Consider the Optimal Control Problem (OCP) associated with a controlled dynamical system:

$$(3-14) \quad \begin{aligned} J^* = \inf_{\mathbf{u}(t)} \int_0^T L(\mathbf{x}(t), \mathbf{u}(t)) dt : & \dot{\mathbf{x}}(t) = f(\mathbf{x}(t), \mathbf{u}(t)), t \in (0, T) \\ & \mathbf{x}(t) \in \mathbf{X}, \mathbf{u}(t) \in \mathbf{U}, \forall t \in (0, T) \\ & \mathbf{x}(0) = \mathbf{x}_0; \mathbf{x}(T) \in \mathbf{X}_T, \end{aligned}$$

where L, f are polynomials, $\mathbf{X}, \mathbf{X}_T \subset \mathbb{R}^n$ and $\mathbf{U} \subset \mathbb{R}^p$ are compact basic semi-algebraic sets. In full generality the OCP problem (Equation (3-14)) is difficult to solve, especially when state constraints $\mathbf{x}(t) \in \mathbf{X}$ are present. Given an admissible state-control trajectory $(t, \mathbf{x}(t), \mathbf{u}(t))$, its associated occupation measure ϕ_1 up to time T (resp. ϕ_2 at time T) are defined by:

$$\phi_1(A \times B \times C) := \int_{[0, T] \cap C} 1_{(A, B)}((\mathbf{x}(t), \mathbf{u}(t))) dt; \quad \phi_2(D) = 1_D(\mathbf{x}(T)),$$

for all $A \in \mathcal{B}(\mathbf{X}), B \in \mathcal{B}(\mathbf{U}), C \in \mathcal{B}([0, T]), D \in \mathcal{B}(\mathbf{X}_T)$. Then for every differentiable function $h : \mathbf{X} \times [0, T] \rightarrow \mathbb{R}$

$$h(T, \mathbf{x}(T)) - h(0, x_0) = \int_0^T \left(\frac{\partial h(\mathbf{x}(t), \mathbf{u}(t))}{\partial t} + \frac{\partial h(\mathbf{x}(t), \mathbf{u}(t))}{\partial \mathbf{x}} f(\mathbf{x}(t), \mathbf{u}(t)) \right) dt,$$

or, equivalently, with $\mathbf{S} := [0, T] \times \mathbf{X} \times \mathbf{U}$:

$$\int_{\mathbf{X}_T} h(T, \mathbf{x}) d\phi_2(\mathbf{x}) = h(0, \mathbf{x}_0) + \int_{\mathbf{S}} \left(\frac{\partial h(\mathbf{x}, \mathbf{u})}{\partial t} + \frac{\partial h(\mathbf{x}, \mathbf{u})}{\partial \mathbf{x}} f(\mathbf{x}, \mathbf{u}) \right) d\phi_1(t, \mathbf{x}, \mathbf{u}).$$

Then *the weak formulation* of the OCP (Equation (3-14)) is the infinite-dimensional linear program:

$$(3-15) \quad \begin{aligned} \rho = \inf_{\phi_1, \phi_2 \geq 0} \{ & \int_{\mathbf{S}} L(\mathbf{x}, \mathbf{u}) d\phi_1 : \\ \text{s.t. } & \int_{\mathbf{X}_T} h(T, \cdot) d\phi_2 - \int_{\mathbf{S}} \left(\frac{\partial h}{\partial t} + \frac{\partial h}{\partial \mathbf{x}} f \right) d\phi_1 = h(0, \mathbf{x}_0) \\ & \forall h \in \mathbb{R}[t, \mathbf{x}] \}. \end{aligned}$$

It turns out that under some conditions the optimal values of Equations (3-14) and (3-15) are equal, i.e., $J^* = \rho$. Next, if one replaces “for all $h \in \mathbb{R}[t, \mathbf{x}, \mathbf{u}]$ ” with “for all $t^k \mathbf{x}^\alpha \mathbf{u}^\beta$ ”, $(t, \alpha, \beta) \in \mathbb{N}^{1+n+p}$ ”, then Equation (3-15) is an instance of the GPM (Equation (3-2)). Therefore one may apply the Moment-SOS hierarchy (Equation (3-5)). Under the conditions of Theorem 3.1 one obtains the asymptotic convergence $\rho_d \rightarrow \rho = J^*$ as $d \rightarrow \infty$. For more details see Lasserre, Henrion, Prieur, and Trélat [2008] and the many references therein.

Robust control. In some applications (e.g. in robust control) one is often interested in optimizing over sets of the form:

$$\mathbf{G} := \{\mathbf{x} \in \Omega_1 : f(\mathbf{x}, \mathbf{u}) \geq 0, \forall \mathbf{u} \in \Omega_2\},$$

where $\Omega_2 \subset \mathbb{R}^p$, and $\Omega_1 \subset \mathbb{R}^n$ is a simple set, in fact a compact set such that one knows all moments of the Lebesgue measure λ on Ω_1 .

The set \mathbf{G} is difficult to handle because of the universal quantifier. Therefore one is often satisfied with an inner approximation $\mathbf{G}_d \subset \mathbf{G}$, and if possible, with (i) a simple form and (ii) some theoretical approximation guarantees. We propose to approximate \mathbf{G} from inside by sets of (simple) form $\mathbf{G}_d = \{\mathbf{x} \in \Omega_1 : p_d(\mathbf{x}) \geq 0\}$ where $p_d \in \mathbb{R}[\mathbf{x}]_{2d}$.

To obtain such an inner approximation $\mathbf{G}_d \subset \mathbf{G}$, define $F : \Omega_1 \rightarrow \mathbb{R}, \mathbf{x} \mapsto F(\mathbf{x}) := \min_{\mathbf{u}} \{f(\mathbf{x}, \mathbf{u}) : \mathbf{u} \in \Omega_2\}$. Then with $d \in \mathbb{N}$, fixed, solve:

$$(3-16) \quad \inf_{p \in \mathbb{R}[\mathbf{x}]_{2d}} \int_{\Omega_1} (F - p) d\lambda : f(\mathbf{x}, \mathbf{u}) - p(\mathbf{x}) \geq 0, \forall (\mathbf{x}, \mathbf{u}) \in \Omega_1 \times \Omega_2\}.$$

Any feasible solution p_d of Equation (3-16) is such that $\mathbf{G}_d = \{\mathbf{x} : p_d(\mathbf{x}) \geq 0\} \subset \mathbf{G}$. In Equation (3-16) $\int_{\Omega_1} (F - p) d\lambda = \|F - p\|_1$ (with $\|\cdot\|_1$ being the $L_1(\Omega_1)$ -norm), and

$$\inf_p \int_{\Omega_1} (F - p) d\lambda = \underbrace{\int_{\Omega_1} F d\lambda}_{=\text{cte}} + \inf_p \int_{\Omega_1} -p d\lambda = \text{cte} - \sup_p \int_{\Omega_1} p d\lambda$$

and so in Equation (3-16) it is equivalent to maximize $\int_{\Omega_1} p d\lambda$. Again the Moment-SOS hierarchy can be applied. This time one replaces the difficult positivity constraint $f(\mathbf{x}, \mathbf{u}) - p(\mathbf{x}) \geq 0$ for all $(\mathbf{x}, \mathbf{u}) \in \Omega_1 \times \Omega_2$ with a certificate of positivity, with a degree bound on the SOS weights. That is, if $\Omega_1 = \{\mathbf{x} : g_{1,\ell}(\mathbf{x}) \geq 0, \ell = 1, \dots, m_1\}$ and $\Omega_2 = \{\mathbf{u} : g_{2,\ell}(\mathbf{u}) \geq 0, \ell = 1, \dots, m_2\}$, then with $d_{i,\ell} := \lceil (\deg(\sigma_{i,\ell})/2) \rceil$, one solves

$$(3-17) \quad \begin{aligned} \rho_d = \sup_{p \in \mathbb{R}[\mathbf{x}]_{2d}} \int_{\Omega_1} p d\lambda : & f(\mathbf{x}, \mathbf{u}) - p(\mathbf{x}) = \sigma_0(\mathbf{x}, \mathbf{u}) \\ & + \sum_{\ell=1}^{m_1} \sigma_{1,\ell}(\mathbf{x}, \mathbf{u}) g_{1,\ell}(\mathbf{x}) + \sum_{\ell=1}^{m_2} \sigma_{2,\ell}(\mathbf{x}, \mathbf{u}) g_{2,\ell}(\mathbf{u}) \\ & \sigma_{i,\ell} \in \Sigma[\mathbf{x}, \mathbf{u}]_{d-d_{i,\ell}}, \ell = 1, \dots, m_i, i = 1, 2. \end{aligned}$$

Theorem 3.2 (Lasserre [2015b]). Assume that $\Omega_1 \times \Omega_2$ is compact and its associated quadratic module is Archimedean. Let p_d be an optimal solution of Equation (3-17). If $\lambda(\{\mathbf{x} \in \Omega_1 : F(\mathbf{x}) = 0\}) = 0$ then $\lim_{d \rightarrow \infty} \|F - p_d\|_1 = 0$ and $\lim_{d \rightarrow \infty} \lambda(\mathbf{G} \setminus \mathbf{G}_d) = 0$.

Therefore one obtains a nested sequence of inner approximations $(\mathbf{G}_d)_{d \in \mathbb{N}} \subset \mathbf{G}$, with the desirable property that $\lambda(\mathbf{G} \setminus \mathbf{G}_d)$ vanishes as d increases. For more details the interested reader is referred to Lasserre [ibid.].

Example 1. In some robust control problems one would like to approximate as closely as desired a non-convex set $\mathbf{G} = \{\mathbf{x} \in \Omega_1 : \lambda_{\min}(\mathbf{A}(\mathbf{x})) \geq 0\}$ for some real symmetric $r \times r$ matrix-polynomial $\mathbf{A}(\mathbf{x})$, and where $\mathbf{x} \mapsto \lambda_{\min}(\mathbf{A}(\mathbf{x}))$ denotes its smallest eigenvalue. If one rewrites

$$\mathbf{G} = \{\mathbf{x} \in \Omega_1 : \mathbf{u}^T \mathbf{A}(\mathbf{x}) \mathbf{u} \geq 0, \forall \mathbf{u} \in \Omega_2\}; \quad \Omega_2 = \{\mathbf{u} \in \mathbb{R}^r : \|\mathbf{u}\| = 1\},$$

one is faced with the problem we have just described. In applying the above methodology the polynomial p_d in [Theorem 3.2](#) approximates $\lambda_{\min}(\mathbf{A}(\mathbf{x}))$ from below in Ω_1 , and $\|p_d(\cdot) - \lambda_{\min}(\mathbf{A}(\cdot))\|_1 \rightarrow 0$ as d increases. For more details see [Henrion and Lasserre \[2006\]](#).

There are many other applications of the Moment-SOS hierarchy in Control, e.g. in Systems Identification [Cerone, Piga, and Regruto \[2012\]](#) and [Laumond, Mansard, and Lasserre \[2017a\]](#), Robotics [Posa, Tobenkin, and Tedrake \[2016\]](#), for computing Lyapunov functions [Parrilo \[2003\]](#), largest regions of attraction [Henrion and Korda \[2014\]](#), to cite a few.

3.6 Some inverse optimization problems. In particular:

Inverse Polynomial Optimization. Here we are given a polynomial optimization problem $\mathbf{P} : f^* = \min\{f(\mathbf{x}) : \mathbf{x} \in \Omega\}$ with $f \in \mathbb{R}[\mathbf{x}]_d$, and we are interested in the following issue: Let $\mathbf{y} \in \Omega$ be given, e.g. \mathbf{y} is the current iterate of a local minimization algorithm applied to \mathbf{P} . Find

$$(3-18) \quad g^* = \arg \min_{g \in \mathbb{R}[\mathbf{x}]_d} \{\|f - g\|_1 : g(\mathbf{x}) - g(\mathbf{y}) \geq 0, \forall \mathbf{x} \in \Omega\},$$

where $\|h\|_1 = \sum_{\alpha} |h_{\alpha}|$ is the ℓ_1 -norm of coefficients of $h \in \mathbb{R}[\mathbf{x}]_d$. In other words, one searches for a polynomial $g^* \in \mathbb{R}[\mathbf{x}]_d$ as close as possible to f and such that $\mathbf{y} \in \Omega$ is a global minimizer of g^* on Ω . Indeed if $\|f - g^*\|_1$ is small enough then $\mathbf{y} \in \Omega$ could be considered a satisfying solution of \mathbf{P} . Therefore given a fixed small $\epsilon > 0$, the test $\|f - g^*\|_1 < \epsilon$ could be a new stopping criterion for a local optimization algorithm, with a strong theoretical justification.

Again the Moment-SOS hierarchy can be applied to solve [Equation \(3-18\)](#) as positivity certificates are perfect tools to handle the positivity constraint “ $g(\mathbf{x}) - g(\mathbf{y}) \geq 0$ for all $\mathbf{x} \in \Omega$ ”. Namely with Ω as in [Equation \(1-2\)](#), solve:

$$(3-19) \quad \rho_t = \min_{g \in \mathbb{R}[\mathbf{x}]_d} \{ \|f - g\|_1 : g(\mathbf{x}) - g(\mathbf{y}) := \sum_{j=0}^m \sigma_j(\mathbf{x}) g_j(\mathbf{x}), \quad \forall \mathbf{x} \},$$

where $g_0(\mathbf{x}) = 1$ for all \mathbf{x} , and $\sigma_j \in \Sigma[\mathbf{x}]_{t-d_j}$, $j = 0, \dots, m$. Other norms are possible but for the sparsity inducing ℓ_1 -norm $\|\cdot\|_1$, it turns out that an optimal solution g^* of [Equation \(3-19\)](#) has a canonical simple form. For more details the interested reader is referred to [Lasserre \[2013\]](#).

Inverse Optimal Control. With the OCP (Equation (3-14)) in Section 3.5, we now consider the following issue: *Given a database of admissible trajectories $(\mathbf{x}(t; \mathbf{x}_\tau), \mathbf{u}(t, \mathbf{x}_\tau))$, $t \in [\tau, T]$, starting in initial state $\mathbf{x}_\tau \in \mathbf{X}$ at time $\tau \in [0, T]$, does there exist a Lagrangian $(\mathbf{x}, \mathbf{u}) \mapsto L(\mathbf{x}, \mathbf{u})$ such that all these trajectories are optimal for the OCP problem (Equation (3-14))?* This problem has important applications, e.g., in Humanoid Robotics to explain human locomotion [Laumond, Mansard, and Lasserre \[2017b\]](#).

Again the Moment-SOS hierarchy can be applied because a weak version of the Hamilton-Jacobi-Bellman (HJB) optimality conditions is the perfect tool to state whether some given trajectory is ϵ -optimal for the OCP (Equation (3-14)). Indeed given $\epsilon > 0$ and an admissible trajectory $(t, \mathbf{x}^*(t), \mathbf{u}^*(t))$, let $\varphi : [0, T] \times \mathbf{X} \rightarrow \mathbb{R}$, and $L : \mathbf{X} \times \mathbf{U} \rightarrow \mathbb{R}$, be such that:

$$(3-20) \quad \varphi(T, \mathbf{x}) \leq 0, \forall \mathbf{x} \in \mathbf{X}; \frac{\partial \varphi(t, \mathbf{x})}{\partial t} + \frac{\partial \varphi(t, \mathbf{x})}{\partial \mathbf{x}} f(\mathbf{x}, \mathbf{u}) + L(\mathbf{x}, \mathbf{u}) \geq 0,$$

for all $(t, \mathbf{x}, \mathbf{u}) \in [0, T] \times \mathbf{X} \times \mathbf{U}$, and: $\varphi(T, \mathbf{x}^*(T)) > -\epsilon$,

$$(3-21) \quad \frac{\partial \varphi(t, \mathbf{x}^*(t))}{\partial t} + \frac{\partial \varphi(t, \mathbf{x}^*(t))}{\partial \mathbf{x}} f(\mathbf{x}^*(t), \mathbf{u}^*(t)) + L(\mathbf{x}^*(t), \mathbf{u}^*(t)) < \epsilon,$$

for all $t \in [0, T]$. Then the trajectory $(t, \mathbf{x}^*(t), \mathbf{u}^*(t))$ is an ϵ -optimal solution of the OCP (Equation (3-14)) with $\mathbf{x}_0 = \mathbf{x}^*(0)$ and Lagrangian L . Therefore to apply the Moment-SOS hierarchy:

(i) The unknown functions φ and L are approximated by polynomials in $\mathbb{R}[t, \mathbf{x}]_{2d}$ and $\mathbb{R}[\mathbf{x}, \mathbf{u}]_{2d}$, where d is the parameter in the Moment-SOS hierarchy (Equation (3-6)).

(ii) The above positivity constraint (Equation (3-20)) on $[0, T] \times \mathbf{X} \times \mathbf{U}$ is replaced with a positivity certificate with degree bound on the SOS weights.

(iii) Equation (3-21) is stated for every trajectory $(\mathbf{x}(t; \mathbf{x}_\tau), \mathbf{u}(t, \mathbf{x}_\tau))$, $t \in [\tau, T]$, in the database. Using a discretization $\{t_1, \dots, t_N\}$ of the interval $[0, T]$, the positivity constraints (Equation (3-21)) then become a set of linear constraints on the coefficients of the unknown polynomials φ and L .

(iv) ϵ in Equation (3-21) is now taken as a variable and one minimizes a criterion of the form $\|L\|_1 + \gamma \epsilon$, where $\gamma > 0$ is chosen to balance between the sparsity-inducing norm $\|L\|_1$ of the Lagrangian and the error ϵ in the weak version of the optimality conditions (Equation (3-20)inv2). A detailed discussion and related results can be found in [Pauwels, Henrion, and Lasserre \[2016\]](#).

3.7 Optimal design in statistics. In designing experiments one models the responses z_1, \dots, z_N of a random *experiment* whose inputs are represented by a vector $\mathbf{t} = (t_i) \in \mathbb{R}^n$ with respect to known *regression functions* $\Phi = (\varphi_1, \dots, \varphi_p)$, namely: $z_i = \sum_{j=1}^p \theta_j \varphi_j(t_i) + \varepsilon_i$, $i = 1, \dots, N$, where $\theta_1, \dots, \theta_p$ are unknown parameters that the experimenter wants to estimate, ε_i is some noise and the (t_i) 's are chosen by the experimenter in a *design space* $\mathcal{X} \subseteq \mathbb{R}^n$. Assume that the inputs t_i , $i = 1, \dots, N$, are chosen within a set of distinct points $\mathbf{x}_1, \dots, \mathbf{x}_\ell \in \mathcal{X}$, $\ell \leq \mathbb{N}$, and let n_k denote the

number of times the particular point \mathbf{x}_k occurs among t_1, \dots, t_N . A design ξ is then defined by:

$$(3-22) \quad \xi = \begin{pmatrix} \mathbf{x}_1 & \dots & \mathbf{x}_\ell \\ \frac{n_1}{N} & \dots & \frac{n_\ell}{N} \end{pmatrix}.$$

The matrix $\mathbf{M}(\xi) := \sum_{i=1}^{\ell} \frac{n_i}{N} \Phi(\mathbf{x}_i) \Phi(\mathbf{x}_i)^T$ is called the information matrix of ξ . Optimal design is concerned with finding a set of points in \mathfrak{X} that optimizes a certain statistical criterion $\phi(\mathbf{M}(\xi))$, which must be real-valued, positively homogeneous, non constant, upper semi-continuous, isotonic w.r.t. Loewner ordering, and concave. For instance in *D-optimal design* one maximizes $\phi(\mathbf{M}(\xi)) := \log \det(\mathbf{M}(\xi))$ over all ξ of the form (Equation (3-22)). This is a difficult problem and so far most methods have used a discretization of the design space \mathfrak{X} .

The Moment-SOS hierarchy that we describe below does not rely on any discretization and works for an arbitrary compact basic semi-algebraic design space \mathfrak{X} as defined in Equation (1-2). Instead we look for an atomic measure on \mathfrak{X} (with finite support) and we proceed in two steps:

- In the first step one solves the hierarchy of convex optimization problems indexed by $\delta = 0, 1, \dots$

$$(3-23) \quad \rho_\delta = \sup_{\mathbf{y}} \{ \log \det(\mathbf{M}_d(\mathbf{y})) : y_0 = 1, \mathbf{M}_{d+\delta}(\mathbf{y}) \succeq 0; \mathbf{M}_{d+\delta-d_j}(g_j \mathbf{y}) \succeq 0 \},$$

where d is fixed by the number of basis functions φ_j considered (here the monomials $(\mathbf{x}^\alpha)_{\alpha \in \mathbb{N}_d^n}$). (Note that Equation (3-23) is not an SDP because the criterion is not linear in \mathbf{y} , but it is still a tractable convex problem.) This provides us with an optimal solution $\mathbf{y}^*(\delta)$. In practice one chooses $\delta = 0$.

- In a second step we extract an atomic measure μ from the “moments” $\mathbf{y}^*(\delta)$, e.g. via Nie’s method Nie [2014b] which consists of solving the SDP:

$$(3-24) \quad \rho_r = \sup_{\mathbf{y}} \{ L_{\mathbf{y}}(f_r) : y_\alpha = y_\alpha^*(\delta), \forall \alpha \in \mathbb{N}_{2d}^n, \mathbf{M}_{d+r}(\mathbf{y}) \succeq 0; \mathbf{M}_{d+r-d_j}(g_j \mathbf{y}) \succeq 0 \},$$

where f_r is a (randomly chosen) polynomial strictly positive on \mathfrak{X} . If $(y_\alpha^*(\delta))_{\alpha \in \mathbb{N}_{2d}^n}$ has a representing measure then it has an atomic representing measure, and generically the rank condition (Equation (2-10)) will be satisfied. Extraction of atoms is obtained via a linear algebra routine. We have tested this two-steps method on several non-trivial numerical experiments (in particular with highly non-convex design spaces \mathfrak{X}) and in all cases we were able to obtain a design. For more details the interested reader is referred to De Castro, Gamboa, Henrion, Hess, and Lasserre [2017].

Other applications & extensions. In this partial overview, by lack of space we have not described some impressive success stories of the Moment-SOS hierarchy, e.g. in coding Bachoc and Vallentin [2008], packing problems in discrete geometry de Laet

and Vallentin [2015] and Schürmann and Vallentin [2006]. Finally, there is also a *non-commutative* version Pironio, Navascués, and Acín [2010] of the Moment-SOS hierarchy based on non-commutative positivity certificates Helton and McCullough [2004] and with important applications in quantum information Navascués, Pironio, and Acín [2008].

4 Conclusion

The list of important applications of the GPM is almost endless and we have tried to convince the reader that the Moment-SOS hierarchy is one promising powerful tool for solving the GPM with already some success stories. However much remains to be done as its brute force application does not scale well to the problem size. One possible research direction is to exploit symmetries and/or sparsity in large scale problems. Another one is to determine alternative positivity certificates which are less expensive in terms of computational burden to avoid the size explosion of SOS-based positivity certificates.

References

- Christine Bachoc and Frank Vallentin (2008). “New upper bounds for kissing numbers from semidefinite programming”. *J. Amer. Math. Soc.* 21.3, pp. 909–924. MR: [2393433](#) (cit. on p. 3778).
- Boaz Barak and David Steurer (2014). “Sum-of-squares proofs and the quest toward optimal algorithms”. In: *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. IV*. Kyung Moon Sa, Seoul, pp. 509–533. MR: [3727623](#) (cit. on p. 3769).
- Emmanuel J. Candès and Carlos Fernandez-Granda (2014). “Towards a mathematical theory of super-resolution”. *Comm. Pure Appl. Math.* 67.6, pp. 906–956. MR: [3193963](#) (cit. on p. 3773).
- Vito Cerone, Dario Piga, and Diego Regruto (2012). “Set-membership error-in-variables identification through convex relaxation techniques”. *IEEE Trans. Automat. Control* 57.2, pp. 517–522. MR: [2918760](#) (cit. on p. 3776).
- Yohann De Castro, F. Gamboa, Didier Henrion, and Jean B. Lasserre (2017). “Exact solutions to super resolution on semi-algebraic domains in higher dimensions”. *IEEE Trans. Inform. Theory* 63.1, pp. 621–630. MR: [3599963](#) (cit. on p. 3773).
- Yohann De Castro, Fabrice Gamboa, Didier Henrion, Roxana Hess, and Jean B. Lasserre (2017). “Approximate Optimal Designs for Multivariate Polynomial Regression”. LAAS report No 17044. 2017, Toulouse, France. To appear in *Annals of Statistics*. arXiv: [1706.04059](#) (cit. on p. 3778).
- Michel X. Goemans and David P. Williamson (1995). “Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming”. *J. Assoc. Comput. Mach.* 42.6, pp. 1115–1145. MR: [1412228](#) (cit. on p. 3762).

- J. William Helton and Scott A. McCullough (2004). “A Positivstellensatz for non-commutative polynomials”. *Trans. Amer. Math. Soc.* 356.9, pp. 3721–3737. MR: 2055751 (cit. on p. 3779).
- D. Henrion, Jean B. Lasserre, and C. Savorgnan (2009). “Approximate volume and integration for basic semialgebraic sets”. *SIAM Rev.* 51.4, pp. 722–743. MR: 2563831 (cit. on p. 3772).
- Didier Henrion and Milan Korda (2014). “Convex computation of the region of attraction of polynomial control systems”. *IEEE Trans. Automat. Control* 59.2, pp. 297–312. MR: 3164876 (cit. on p. 3776).
- Didier Henrion and Jean B. Lasserre (2006). “Convergent relaxations of polynomial matrix inequalities and static output feedback”. *IEEE Trans. Automat. Control* 51.2, pp. 192–202. MR: 2201707 (cit. on p. 3776).
- Didier Henrion, Jean B. Lasserre, and Johan Löfberg (2009). “GloptiPoly 3: moments, optimization and semidefinite programming”. *Optim. Methods Softw.* 24.4–5, pp. 761–779. MR: 2554910 (cit. on p. 3767).
- Cédric Jozs, Jean B. Lasserre, and Bernard Mourrain (Aug. 2017). “Sparse polynomial interpolation: compressed sensing, super resolution, or Prony?” LAAS Report no 17279. 2017, Toulouse, France. arXiv: 1708.06187 (cit. on pp. 3773, 3774).
- Subhash Khot (2010). “Inapproximability of NP-complete problems, discrete Fourier analysis, and geometry”. In: *Proceedings of the International Congress of Mathematicians. Volume IV*. Hindustan Book Agency, New Delhi, pp. 2676–2697. MR: 2827989 (cit. on p. 3769).
- (2014). “Hardness of approximation”. In: *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. 1*. Kyung Moon Sa, Seoul, pp. 711–728. MR: 3728489 (cit. on pp. 3762, 3769).
- Etienne de Klerk, Jean B. Lasserre, Monique Laurent, and Zhao Sun (2017). “Bound-constrained polynomial optimization using only elementary calculations”. *Math. Oper. Res.* 42.3, pp. 834–853. MR: 3685268.
- J.-L. Krivine (1964a). “Anneaux préordonnés”. *J. Analyse Math.* 12, pp. 307–326. MR: 0175937 (cit. on p. 3765).
- Jean-Louis Krivine (1964b). “Quelques propriétés des préordres dans les anneaux commutatifs unitaires”. *C. R. Acad. Sci. Paris* 258, pp. 3417–3418. MR: 0169083 (cit. on p. 3765).
- David de Laat and Frank Vallentin (2015). “A semidefinite programming hierarchy for packing problems in discrete geometry”. *Math. Program.* 151.2, Ser. B, pp. 529–553. MR: 3348162 (cit. on p. 3778).
- Henry J Landau (1987). *Moments in mathematics*. Vol. 37. Proc. Sympos. Appl. Math. (cit. on p. 3763).
- Jean B. Lasserre (2000). “Optimisation globale et théorie des moments”. *C. R. Acad. Sci. Paris Sér. I Math.* 331.11, pp. 929–934. MR: 1806434 (cit. on pp. 3762, 3767).
- (2002a). “Bounds on measures satisfying moment conditions”. *Ann. Appl. Probab.* 12.3, pp. 1114–1137. MR: 1925454 (cit. on p. 3771).

- Jean B. Lasserre (2002b). “Semidefinite programming vs. LP relaxations for polynomial programming”. *Math. Oper. Res.* 27.2, pp. 347–360. MR: 1908532 (cit. on pp. 3766, 3769).
- (2006). “Convergent SDP-relaxations in polynomial optimization with sparsity”. *SIAM J. Optim.* 17.3, pp. 822–843. MR: 2257211 (cit. on p. 3768).
 - (2010). *Moments, positive polynomials and their applications*. Vol. 1. Imperial College Press Optimization Series. Imperial College Press, London, pp. xxii+361. MR: 2589247 (cit. on p. 3765).
 - (2013). “Inverse polynomial optimization”. *Math. Oper. Res.* 38.3, pp. 418–436. MR: 3092539 (cit. on p. 3776).
 - (2015a). *An introduction to polynomial and semi-algebraic optimization*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, pp. xiv+339. MR: 3469431 (cit. on pp. 3766, 3768, 3769).
 - (2015b). “Tractable approximations of sets defined with quantifiers”. *Math. Program.* 151.2, Ser. B, pp. 507–527. MR: 3348161 (cit. on p. 3775).
 - (2017). “Computing Gaussian & exponential measures of semi-algebraic sets”. *Adv. in Appl. Math.* 91, pp. 137–163. MR: 3673583 (cit. on p. 3772).
 - (2000/01). “Global optimization with polynomials and the problem of moments”. *SIAM J. Optim.* 11.3, pp. 796–817. MR: 1814045 (cit. on pp. 3762, 3767).
- Jean B. Lasserre, Didier Henrion, Christophe Prieur, and Emmanuel Trélat (2008). “Nonlinear optimal control via occupation measures and LMI-relaxations”. *SIAM J. Control Optim.* 47.4, pp. 1643–1666. MR: 2421324 (cit. on p. 3774).
- Jean B. Lasserre, Monique Laurent, and Philipp Rostalski (2008). “Semidefinite characterization and computation of zero-dimensional real radical ideals”. *Found. Comput. Math.* 8.5, pp. 607–647. MR: 2443091.
- Jean-Paul Laumond, Nicolas Mansard, and Jean B. Lasserre, eds. (2017a). *Geometric and numerical foundations of movements*. Vol. 117. Springer Tracts in Advanced Robotics. Springer, Cham, pp. x+419. MR: 3642945 (cit. on pp. 3768, 3776).
- eds. (2017b). *Geometric and numerical foundations of movements*. Vol. 117. Springer Tracts in Advanced Robotics. Springer, Cham, pp. x+419. MR: 3642945 (cit. on p. 3777).
- Monique Laurent (2003). “A comparison of the Sherali-Adams, Lovász-Schrijver, and Lasserre relaxations for 0-1 programming”. *Math. Oper. Res.* 28.3, pp. 470–496. MR: 1997246 (cit. on p. 3769).
- (2009). “Sums of squares, moment matrices and optimization over polynomials”. In: *Emerging applications of algebraic geometry*. Vol. 149. IMA Vol. Math. Appl. Springer, New York, pp. 157–270. MR: 2500468 (cit. on p. 3765).
- Daniel K Molzahn and Ian A Hiskens (2015). “Sparsity-exploiting moment-based relaxations of the optimal power flow problem”. *IEEE Transactions on Power Systems* 30.6, pp. 3168–3180 (cit. on p. 3768).
- Miguel Navascués, Stefano Pironio, and Antonio Acín (2008). “A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations”. *New Journal of Physics* 10.7, p. 073013 (cit. on p. 3779).

- Yurii Nesterov (2000). “Squared functional systems and optimization problems”. In: *High performance optimization*. Vol. 33. Appl. Optim. Kluwer Acad. Publ., Dordrecht, pp. 405–440. MR: [1748764](#) (cit. on p. [3762](#)).
- Jiawang Nie (2013). “Certifying convergence of Lasserre’s hierarchy via flat truncation”. *Math. Program.* 142.1-2, Ser. A, pp. 485–510. MR: [3127083](#) (cit. on pp. [3768](#), [3773](#)).
- (2014a). “Optimality conditions and finite convergence of Lasserre’s hierarchy”. *Math. Program.* 146.1-2, Ser. A, pp. 97–121. MR: [3232610](#) (cit. on pp. [3767](#), [3768](#)).
- (2014b). “The \mathcal{A} -truncated K -moment problem”. *Found. Comput. Math.* 14.6, pp. 1243–1276. MR: [3273678](#) (cit. on p. [3778](#)).
- Pablo A. Parrilo (2000). “Structured Semidefinite Programs and Semialgebraic Geometry Methods in Robustness and Optimization”. PhD thesis. California Institute of Technology (cit. on p. [3762](#)).
- (2003). “Semidefinite programming relaxations for semialgebraic problems”. *Math. Program.* 96.2, Ser. B. Algebraic and geometric methods in discrete optimization, pp. 293–320. MR: [1993050](#) (cit. on pp. [3762](#), [3776](#)).
- Edouard Pauwels, Didier Henrion, and Jean B. Lasserre (2016). “Linear conic optimization for inverse optimal control”. *SIAM J. Control Optim.* 54.3, pp. 1798–1825. MR: [3516862](#) (cit. on p. [3777](#)).
- S. Pironio, M. Navascués, and A. Acín (2010). “Convergent relaxations of polynomial optimization problems with noncommuting variables”. *SIAM J. Optim.* 20.5, pp. 2157–2180. MR: [2650843](#) (cit. on p. [3779](#)).
- Michael Posa, Mark Tobenkin, and Russ Tedrake (2016). “Stability analysis and control of rigid-body systems with impacts and friction”. *IEEE Trans. Automat. Control* 61.6, pp. 1423–1437. MR: [3508689](#) (cit. on p. [3776](#)).
- Mihai Putinar (1993). “Positive polynomials on compact semi-algebraic sets”. *Indiana Univ. Math. J.* 42.3, pp. 969–984. MR: [1254128](#) (cit. on p. [3765](#)).
- Konrad Schmüdgen (2017). *The moment problem*. Vol. 277. Graduate Texts in Mathematics. Springer, Cham, pp. xii+535. MR: [3729411](#) (cit. on p. [3765](#)).
- Achill Schürmann and Frank Vallentin (2006). “Computational approaches to lattice packing and covering problems”. *Discrete Comput. Geom.* 35.1, pp. 73–116. MR: [2183491](#) (cit. on pp. [3778](#), [3779](#)).
- Naum Z. Shor (1998). *Nondifferentiable optimization and polynomial problems*. Vol. 24. Nonconvex Optimization and its Applications. Kluwer Academic Publishers, Dordrecht, pp. xviii+394. MR: [1620179](#) (cit. on p. [3762](#)).
- F.-H. Vasilescu (2003). “Spectral measures and moment problems”. In: *Spectral analysis and its applications*. Vol. 2. Theta Ser. Adv. Math. Theta, Bucharest, pp. 173–215. MR: [2082433](#) (cit. on p. [3765](#)).
- Hayato Waki, Sunyoung Kim, Masakazu Kojima, and Masakazu Muramatsu (2006). “Sums of squares and semidefinite program relaxations for polynomial optimization problems with structured sparsity”. *SIAM J. Optim.* 17.1, pp. 218–242. MR: [2219151](#) (cit. on p. [3768](#)).

JEAN B. LASSERRE
lasserre@laas.fr

